Multiple Quantum Hypothesis Testing Expressions and Classical-Quantum Channel Converse Bounds

Gonzalo Vazquez-Vilar
Universidad Carlos III de Madrid, Spain
Email: gvazquez@ieee.org

arXiv:1607.07625v1 [quant-ph] 26 Jul 2016

***Abstract*—Alternative exact expressions are derived for the minimum error probability of a hypothesis test discriminating among M quantum states. The first expression corresponds to the error probability of a binary hypothesis test with certain parameters; the second involves the optimization of a given information-spectrum measure. Particularized in the classical-quantum channel coding setting, this characterization implies the tightness of two existing converse bounds; one derived by Matthews and Wehner using hypothesis-testing, and one obtained by Hayashi and Nagaoka via an information-spectrum approach.**

I. Introduction

Optimal discrimination among multiple quantum states –quantum hypothesis testing– is at the core of several information processing tasks involving quantum-mechanical systems. When the number of hypotheses is two, quantum hypothesis testing allows a simple formulation in terms of two kinds of pairwise errors. The quantum version of the Neyman-Pearson lemma establishes the optimum binary test in this setting. This problem was first studied by Helstrom in [1] (see also [2], [3]). When the number of hypotheses is larger than two, a (classical) prior distribution is usually placed over the hypotheses. While there exists no closed form for the optimal test in general, optimality contitions can be obtained [4], [5]. For historical notes on the subject see [6, Ch. IV].

In the context of reliable communication, hypothesis testing has been instrumental in the derivation of converse bounds to the error probability both in the classical and quantum settings (see, e.g., [7], [8]). Recently, hypothesis testing gained interest as a very general approach to obtain converse bounds in the finite block-length regime. In classical channel coding, Polyanskiy, Poor and Verdú derived the meta-converse bound based on an instance of binary hypothesis testing [9]. A similar approach was used by Wang and Renner to derive a finite block-length converse bound for classical-quantum channels [10], and by Matthews and Wehner to obtain a family of converse bounds for general quantum channels [11]. The results by Matthews and Wehner are general enough to recover the meta-converse bound in the classical setting and Wang-Renner converse bound in the classical-quantum setting.

G. Vazquez-Vilar is also with the Gregorio Marañón Health Research Institute, Madrid, Spain. This work has been funded in part by the Spanish Ministry of Economy and Competitiveness under Grants FPDI-2013-18602, TEC2013-41718-R, and TEC2015-69648-REDC.

The information-spectrum method studies the asymptotics of a certain random variable, often referred to as information density or information random variable. Using a quantum analogue of this quantity, Hayashi and Nagaoka studied quantum hypothesis testing [12], and classical-quantum channel coding [13], obtaining general bounds for both problems.

In this paper, we derive two alternative exact expressions for the minimum error probability of multiple quantum hypothesis testing when a (classical) prior distribution is placed over the hypotheses. The expressions obtained illustrate connections among hypothesis testing, information-spectrum measures and converse bounds in classical-quantum channel coding. An application to classical-quantum channel coding shows that Matthews-Wehner converse bound [11, Th. 19] and Hayashi-Nagaoka lemma [13, Lemma 4] with certain parameters yield the exact error probability. This work thus generalizes several results derived in [14] in the classical setting.

II. Background

A. Notation

In the general case, a quantum state is described by a density operator ρ acting on some finite dimensional complex Hilbert space $\mathcal{H}$. Density operators are self-adjoint, positive semidefinite, and have unit trace. A measurement on a quantum system is a mapping from the state of the system ρ to a classical outcome $m \in \{1, \dots, M\}$. A measurement is represented by a collection of positive self-adjoint operators $\{\Pi_1, \dots, \Pi_M\}$ such that $\sum \Pi_m = \mathbb{1}$, where $\mathbb{1}$ is the identity operator. These operators form a POVM (positive operator-valued measure). A measurement $\{\Pi_1, \dots, \Pi_M\}$ applied to ρ has outcome m with probability $\mathrm{Tr}(\rho\Pi_m)$.

For two self-adjoint operators A, B, the notation $A \geq B$ means that $A - B$ is positive semidefinite. Similarly $A \leq B$, $A > B$, and $A < B$ means that $A - B$ is negative semidefinite, positive definite and negative definite, respectively. For a self-adjoint operator A with spectral decomposition $A = \sum_i \lambda_i E_i$, where $\{\lambda_i\}$ are the eigenvalues and $\{E_i\}$ are the orthogonal projections onto the corresponding eigenspaces, we define

$$\{A > 0\} \triangleq \sum_{i:\lambda_i > 0} E_i. \tag{1}$$

This corresponds to the projector associated to the positive eigenspace of A. We shall also use $\{A \geq 0\} \triangleq \sum_{i:\lambda_i \geq 0} E_i$, $\{A < 0\} \triangleq \sum_{i:\lambda_i < 0} E_i$ and $\{A \leq 0\} \triangleq \sum_{i:\lambda_i \leq 0} E_i$.

B. Binary Hypothesis Testing

Let us consider a binary hypothesis test (with simple hypotheses) discriminating between the density operators ρ_0 and ρ_1 acting on $\mathcal{H}$. In order to distinguish between the two hypotheses we perform a measurement. We define a test measurement $\{T, \bar{T}\}$, such that T and $\bar{T} \triangleq \mathbb{1} - T$ are positive semidefinite. The test decides ρ_0 (resp. ρ_1) when the measurement outcome corresponding to T (resp. $\bar{T}$) occurs.

Let $\epsilon_{j|i}$ denote the probability of deciding ρ_j when ρ_i is the true hypothesis, $i, j = 0, 1$, $i \neq j$. More precisely, we define

$$\epsilon_{1|0}(T) \triangleq 1 - \operatorname{Tr}(\rho_0 T) = \operatorname{Tr}\left(\rho_0 \bar{T}\right), \tag{2}$$

$$\epsilon_{0|1}(T) \triangleq \operatorname{Tr}(\rho_1 T). \tag{3}$$

Let $\alpha_\beta(\rho_0 \| \rho_1)$ denote the minimum error probability $\epsilon_{1|0}$ among all tests with $\epsilon_{0|1}$ at most β, that is,

$$\alpha_\beta(\rho_0 \| \rho_1) \triangleq \inf_{T : \epsilon_{0|1}(T) \leq \beta} \epsilon_{1|0}(T). \tag{4}$$

The function $\alpha_\beta(\cdot\|\cdot)$ is the inverse of the function $\beta_\alpha(\cdot\|\cdot)$ appearing in [11], which is itself related to the hypothesis-testing relative entropy as $D_{\mathrm{H}}^{\alpha}(\rho_0\|\rho_1) = -\log \beta_\alpha(\rho_0\|\rho_1)$ [10].

When ρ_0 and ρ_1 commute, the test T in (4) can be restricted to be diagonal in the (common) eigenbasis of ρ_0 and ρ_1, then (4) reduces to the classical case [14].

The quantum version of the Neyman-Pearson lemma characterizes the form of the test minimizing (4). Let $t \geq 0$ and let P_t^+, P_t^-, P_t^0 denote the projectors spanning the positive, negative and null eigenspaces of the matrix $\rho_0 - t\rho_1$, respectively, i. e.,

$$P_t^+ \triangleq \left\{\rho_0 - t\rho_1 > 0\right\}, \tag{5}$$

$$P_t^- \triangleq \left\{\rho_0 - t\rho_1 < 0\right\}, \tag{6}$$

$$P_t^0 \triangleq \mathbb{1} - P_t^+ - P_t^-. \tag{7}$$

Lemma 1 (Neyman-Pearson lemma): The operator T_{NP} is an optimal test between ρ_0 and ρ_1 if and only if

$$T_{\mathrm{NP}} = P_t^+ + p_t^0, \tag{8}$$

where $0 \leq p_t^0 \leq P_t^0$.

Proof: A slightly different formulation of this result is usually given in the literature. The statement included here can be found in, e.g., [15, Lem. 3]. ∎

Therefore, for any $t \geq 0$ and $0 \leq p_t^0 \leq P_t^0$ such that $\operatorname{Tr}\{\rho_1 T_{\mathrm{NP}}\} = \beta$, the resulting test T_{NP} minimizes (4). Moreover, the following lower bound holds.

Lemma 2: For any test discriminating between ρ_0 and ρ_1, and for any $t' \geq 0$, it holds that

$$\alpha_\beta(\rho_0\|\rho_1) \;\geq\; \operatorname{Tr}\Big(\rho_0\big(P_{t'}^- + P_{t'}^0\big)\Big) - t'\beta. \tag{9}$$

Proof: For any operator $A \geq 0$ and $0 \leq T \leq \mathbb{1}$, it holds that $\operatorname{Tr}\big(A\{A > 0\}\big) \geq \operatorname{Tr}(AT)$ [12, Eq. 8]. For $A = \rho_0 - t'\rho_1$ and $T = T_{\mathrm{NP}}$, this inequality becomes

$$\operatorname{Tr}\big((\rho_0 - t'\rho_1)P_{t'}^+\big) \geq \operatorname{Tr}\big((\rho_0 - t'\rho_1)T_{\mathrm{NP}}\big), \tag{10}$$

which after some algebra yields

$$-\operatorname{Tr}\big(\rho_0 T_{\mathrm{NP}}\big) \geq -\operatorname{Tr}\big(\rho_0 P_{t'}^+\big) + t' \operatorname{Tr}\big(\rho_1(P_{t'}^+ - T_{\mathrm{NP}})\big). \tag{11}$$

Summing one to both sides of (11) and noting that $\alpha_\beta(\rho_0\|\rho_1) = 1 - \operatorname{Tr}\big(\rho_0 T_{\mathrm{NP}}\big)$ and $\beta = \operatorname{Tr}\big(\rho_1 T_{\mathrm{NP}}\big)$, we obtain

$$\alpha_\beta(\rho_0\|\rho_1) \geq \operatorname{Tr}\big(\rho_0(P_{t'}^- + P_{t'}^0)\big) + t'\operatorname{Tr}\big(\rho_1 P_{t'}^+\big) - t'\beta. \tag{12}$$

The result thus follows by lower-bounding $\operatorname{Tr}\big(\rho_1 P_{t'}^+\big) \geq 0$. ∎

III. Multiple Quantum Hypothesis Testing

We consider a hypothesis testing problem discriminating among M possible states acting on $\mathcal{H}$, where M is assumed to be finite. The M alternatives $\tau_1, \ldots, \tau_M$ are assumed to occur with (classical) probabilities $p_1, \ldots, p_M$, respectively.

A M-ary hypothesis test is a POVM $\mathcal{P} \triangleq \{\Pi_1, \Pi_2, \ldots, \Pi_M\}$, $\sum \Pi_i = \mathbb{1}$. The test decides the alternative τ_i when the measurement with respect to $\mathcal{P}$ has outcome i. The probability that the test $\mathcal{P}$ decides τ_j when τ_i is the true underlying state is thus $\operatorname{Tr}\big(\tau_i \Pi_j\big)$ and the average error probability is

$$\epsilon(\mathcal{P}) \triangleq 1 - \sum_{i=1}^{M} p_i \operatorname{Tr}(\tau_i \Pi_i). \tag{13}$$

We define the minimum average error probability as

$$\epsilon \triangleq \min_{\mathcal{P}} \epsilon(\mathcal{P}). \tag{14}$$

The test $\mathcal{P}$ minimizing (14) has no simple form in general.

Lemma 3 (Holevo-Yuen-Kennedy-Lax conditions): A test $\mathcal{P}^\star = \{\Pi_1^\star, \ldots, \Pi_M^\star\}$ minimizes (14) if and only if, for each $m = 1, \ldots, M$,

$$\big(\Lambda(\mathcal{P}^\star) - p_m\tau_m\big)\Pi_m^\star \;=\; \Pi_m^\star\big(\Lambda(\mathcal{P}^\star) - p_m\tau_m\big) \;=\; 0, \tag{15}$$

$$\Lambda(\mathcal{P}^\star) - p_m\tau_m \;\geq\; 0, \tag{16}$$

where

$$\Lambda(\mathcal{P}^\star) \triangleq \sum_{i=1}^{M} p_i\tau_i\Pi_i^\star = \sum_{i=1}^{M} p_i\Pi_i^\star\tau_i \tag{17}$$

is required to be self-adjoint[1].

Proof: The theorem follows from [4, Th. 4.1, Eq. (4.8)] or [5, Th. I] after simplifying the resulting optimality conditions. ∎

We next show an alternative characterization of the minimum error probability ϵ as a function of a binary hypothesis test with certain parameters.

Let $\operatorname{diag}(\rho_1, \ldots, \rho_M)$ denote the block-diagonal matrix with diagonal blocks $\rho_1, \ldots, \rho_M$. We define

$$\mathcal{T} \triangleq \operatorname{diag}\big(p_1\tau_1, \ldots, p_M\tau_M\big), \tag{18}$$

$$\mathcal{D}(\mu_0) \triangleq \operatorname{diag}\big(\tfrac{1}{M}\mu_0, \ldots, \tfrac{1}{M}\mu_0\big), \tag{19}$$

where μ_0 is an arbitrary density operator acting on $\mathcal{H}$. Note that both $\mathcal{T}$ and $\mathcal{D}(\mu_0)$ are density operators themselves, since they are self-adjoint, positive semidefinite and have unit trace.

[1]The operator $\Lambda(\mathcal{P})$ takes a role of the Lagrange multiplier associated to the constraint $\sum \Pi_i = \mathbb{1}$, which, involving self-adjoint operators requires Λ to be self-adjoint.

Theorem 1: The minimum error probability of an M-ary test discriminating among states $\{\tau_1, \ldots, \tau_M\}$ with prior classical probabilities $\{p_1, \ldots, p_M\}$ satisfies

$$\epsilon = \max_{\mu_0} \alpha_{\frac{1}{M}}\big(\mathcal{T} \,\|\, \mathcal{D}(\mu_0)\big), \tag{20}$$

where $\mathcal{T}$ and $\mathcal{D}(\cdot)$ are given in (18) and (19), respectively, and where the optimization is carried out over (unit-trace non-negative) density operators μ_0.

Proof: For any $\mathcal{P} = \{\Pi_1, \Pi_2, \ldots, \Pi_M\}$ let us define the binary test $T' \triangleq \operatorname{diag}(\Pi_1, \ldots, \Pi_M)$. For this test we obtain

$$\epsilon_{1|0}(T') = 1 - \sum_{i=1}^{M} p_i \operatorname{Tr}(\tau_i \Pi_i) = \epsilon(\mathcal{P}), \tag{21}$$

$$\epsilon_{0|1}(T') = \frac{1}{M} \sum_{i=1}^{M} \operatorname{Tr}(\mu_0 \Pi_i) \tag{22}$$

$$= \frac{1}{M} \operatorname{Tr}\left(\mu_0 \left(\sum\nolimits_{i=1}^{M} \Pi_i\right)\right) \tag{23}$$

$$= \frac{1}{M} \operatorname{Tr}(\mu_0) = \frac{1}{M}. \tag{24}$$

The (possibly suboptimal) test T' has thus $\epsilon_{1|0}(T') = \epsilon(\mathcal{P})$ and $\epsilon_{0|1}(T') = \frac{1}{M}$. Therefore, using (4) and maximizing the resulting expression over μ_0, we obtain

$$\epsilon(\mathcal{P}) \geq \max_{\mu_0} \alpha_{\frac{1}{M}}\big(\mathcal{T} \,\|\, \mathcal{D}(\mu_0)\big). \tag{25}$$

It remains to show that, for $\mathcal{P} = \mathcal{P}^\star$ defined in Lemma 3, the lower bound (25) holds with equality. To this end, we next demonstrate that the optimality conditions for T_{NP} in Lemma 1 and for $\mathcal{P}^\star = \{\Pi_1^\star, \ldots, \Pi_M^\star\}$ in Lemma 3 are equivalent for a specific choice of μ_0.

Let $\mathcal{P}^\star = \{\Pi_1^\star, \ldots, \Pi_M^\star\}$ satisfy (15)-(16) and define

$$\mu_0^\star \triangleq \frac{1}{c_0^\star} \sum_{i=1}^{M} p_i \tau_i \Pi_i^\star = \frac{1}{c_0^\star} \Lambda(\mathcal{P}^\star), \tag{26}$$

where $c_0^\star$ is a normalizing constant such that $\mu_0^\star$ is unit trace.

Lemma 1 shows that the test T_{NP} achieving (25) is associated to the non-negative eigenspace of the matrix $\mathcal{T} - t\mathcal{D}(\mu_0)$. Given the block-diagonal structure of the matrix $\mathcal{T} - t\mathcal{D}(\mu_0)$, it is enough to consider binary tests T_{NP} with block-diagonal structure. Then, we write $T_{\text{NP}} = \operatorname{diag}\big(T_1^{\text{NP}}, \ldots, T_M^{\text{NP}}\big)$.

For the choice $\mu_0 = \mu_0^\star$, and $t = Mc_0^\star$, the m-th block-diagonal term in $\mathcal{T} - t\mathcal{D}(\mu_0)$ is given by

$$p_m \tau_m - \tfrac{t}{M} \mu_0 = p_m \tau_m - \Lambda(\mathcal{P}^\star). \tag{27}$$

The m-th block of the Neyman-Pearson test T_m^{NP} must lie in the non-negative eigenspace of the matrix (27). However, since (16) implies that (27) is negative semidefinite, each block T_m^{NP} can only lie in the null eigenspace of (27), $m = 1, \ldots, M$.

According to (15), the operator $\Pi_m^\star$ belongs to the null eigenspace of (27), $m = 1, \ldots, M$. As a result, the choice

$$T_{\text{NP}} = \operatorname{diag}(\Pi_1^\star, \ldots, \Pi_M^\star) \tag{28}$$

satisfies the optimality conditions in Lemma 1. Moreover, since $\epsilon_{1|0}(T_{\text{NP}}) = \epsilon\big(\mathcal{P}^\star\big) = \epsilon$ and $\epsilon_{0|1}(T_{\text{NP}}) = \frac{1}{M}$, Lemma 1 implies that (20) holds with equality for $\mu_0 = \mu_0^\star$. Given the bound in (25), other choices of μ_0 cannot improve the result, and Theorem 1 thus follows. ∎

Combining Theorem 1 and Lemma 2, we obtain a characterization for ϵ based on information-spectrum measures.

Theorem 2: The minimum error probability of an M-ary test discriminating among states $\{\tau_1, \ldots, \tau_M\}$ with prior classical probabilities $\{p_1, \ldots, p_M\}$ satisfies

$$\epsilon = \max_{\mu_0, t \geq 0} \left\{ \sum_{i=1}^{M} p_i \operatorname{Tr}\Big(\tau_i \big\{p_i \tau_i - t\mu_0 \leq 0\big\}\Big) - t \right\}. \tag{29}$$

where the optimization is carried out over (unit-trace non-negative) density operators μ_0 acting on $\mathcal{H}$, and over the scalar threshold $t \geq 0$.

Proof: Applying Lemma 2 to (20), and using the definitions of $\mathcal{T}$ in (18) and $\mathcal{D}(\cdot)$ in (19), yields, for any μ_0, $t' \geq 0$,

$$\epsilon \geq \sum_{i=1}^{M} p_i \operatorname{Tr}\Big(\tau_i \big\{p_i \tau_i - \tfrac{t'}{M} \mu_0 \leq 0\big\}\Big) - \tfrac{t'}{M}. \tag{30}$$

It remains to show that there exist μ_0 and $t' \geq 0$ such that (30) holds with equality. In particular, let us choose $\mu_0 = \mu_0^\star$ defined in (26), and $t' = Mc_0^\star$ where $c_0^\star = \sum_{i=1}^{M} p_i \operatorname{Tr}(\tau_i \Pi_i^\star)$ is the normalizing constant from (26).

For this choice of μ_0 and t', the projector spanning the negative semidefinite eigenspace of the operator $p_i\tau_i - \frac{t'}{M}\mu_0$ can be rewritten as

$$\Big\{p_i \tau_i - \tfrac{t'}{M} \mu_0 \leq 0\Big\} = \big\{p_i \tau_i - \Lambda(\mathcal{P}^\star) \leq 0\big\} \tag{31}$$

$$= \mathbb{1}, \tag{32}$$

where the last identity follows from (16). The right-hand side of (30) thus becomes

$$\sum_{i=1}^{M} p_i \operatorname{Tr}(\tau_i) - \frac{t'}{M} = 1 - \frac{t'}{M}. \tag{33}$$

The result follows since $\frac{t'}{M} = c_0^\star = \sum_i p_i \operatorname{Tr}(\tau_i \Pi_i^\star) = 1 - \epsilon$. ∎

The alternative expressions derived in Theorems 1 and 2 are not easier to compute than the original optimization in (14), all of them requiring to solve a semidefinite program. We recall from the proofs of the theorems that a density operator μ_0 maximizing (20) and (29) is

$$\mu_0^\star = \frac{1}{c_0^\star} \sum_{i=1}^{M} p_i \tau_i \Pi_i^\star, \tag{34}$$

for some $\mathcal{P}^\star = \{\Pi_1^\star, \ldots, \Pi_M^\star\}$ satisfying the conditions in Lemma 3 and where $c_0^\star$ is a normalizing constant. Hence, the optimal M-ary hypothesis test $\mathcal{P}^\star$ characterizes the optimal μ_0. Conversely, the optimal μ_0 is precisely the Lagrange multiplier associated to the minimization in (14), after an appropriate re-scaling.

The expressions derived in Theorems 1 and 2 can be used to determine the tightness of several converse bounds from the literature, as we show in the next section.

IV. Application to Classical-Quantum Channels

We consider the channel coding problem of transmitting M equiprobable messages over a one-shot classical-quantum channel $x \to W_x$, with $x \in \mathcal{X}$ and $W_x \in \mathcal{H}$.

A channel code is defined as a mapping from the message set $\{1, \ldots, M\}$ into a set of M codewords $\mathcal{C} = \{x_1, \ldots, x_M\}$. For a source message m, the decoder receives the associated density operator W_{x_m} and must decide on the transmitted message. The minimum error probability for a code $\mathcal{C}$ is

$$\mathsf{P}_{\mathrm{e}}(\mathcal{C}) \triangleq \min_{\{\Pi_1,\ldots,\Pi_M\}} \left\{ 1 - \frac{1}{M} \sum_{m=1}^{M} \mathrm{Tr}\big(W_{x_m}\Pi_m\big) \right\}. \quad (35)$$

This problem corresponds precisely to the M-ary quantum hypothesis testing problem described in Section III. Then, direct application of Theorems 1 and 2 yields two alternative expressions for $\mathsf{P}_{\mathrm{e}}(\mathcal{C})$.

Let $\mathbb{A}$ and $\mathbb{B}$ denote the input and output of the system, respectively. The joint state induced by a codebook $\mathcal{C}$ is

$$\rho_{\mathcal{C}}^{\mathbb{A}\mathbb{B}} = \frac{1}{M} \sum_{x \in \mathcal{C}} |x\rangle\langle x|^{\mathbb{A}} \otimes W_x^{\mathbb{B}}, \quad (36)$$

and $\rho_{\mathcal{C}}^{\mathbb{A}} = \frac{1}{M} \sum_{x \in \mathcal{C}} |x\rangle\langle x|^{\mathbb{A}}$ its input marginal.

According to (20) in Theorem 1 we obtain

$$\mathsf{P}_{\mathrm{e}}(\mathcal{C}) = \max_{\mu_0} \alpha_{\frac{1}{M}}\big(\rho_{\mathcal{C}}^{\mathbb{A}\mathbb{B}} \,\|\, \rho_{\mathcal{C}}^{\mathbb{A}} \otimes \mu_0^{\mathbb{B}}\big). \quad (37)$$

The expression (37) is precisely the finite block-length converse bound by Matthews and Wehner [11, Eq. (45)], particularized for a classical-quantum channel with an input state induced by the codebook $\mathcal{C}$. Therefore, Theorem 1 implies that the quantum generalization of the meta-converse bound proposed by Matthews and Wehner is tight for a fixed codebook $\mathcal{C}$.

Minimizing the right-hand side of (37) over all distributions P_X defined over the input alphabet $\mathcal{X}$, not necessarily induced by a codebook, yields a lower bound on $\mathsf{P}_{\mathrm{e}}(\mathcal{C})$ for any codebook $\mathcal{C}$. By fixing μ_0 to be the state induced at the system output, this lower bound recovers the converse bound by Wang and Renner [10, Th. 1]. This bound is not tight in general since (i) the minimizing P_X does not need to coincide with the input state induced by the best codebook, and (ii) the choice of μ_0 in [10, Th. 1] does not maximize the resulting bound in general.

Using the characterization in Theorem 2, the error probability $\mathsf{P}_{\mathrm{e}}(\mathcal{C})$ can be equivalently written as

$$\mathsf{P}_{\mathrm{e}}(\mathcal{C}) = \max_{\mu_0, t' \geq 0} \left\{ \frac{1}{M} \sum_{x \in \mathcal{C}} \mathrm{Tr}\Big(W_x\big\{W_x - t'\mu_0 \leq 0\big\}\Big) - \frac{t'}{M} \right\}. \quad (38)$$

The objective of the maximization in (38) coincides with the information-spectrum bound [13, Lemma 4]. Then, (38) shows that the Hayashi-Nagaoka lemma yields the exact error probability for a fixed code, after optimizantion over the free parameters μ_0, $t' \geq 0$.

V. Concluding Remarks

In Theorem 1, the minimum error probability of an M-ary quantum hypothesis test is expressed as an instance of a binary quantum hypothesis test with certain parameters. This expression implies the tightness of the converse bound [11, Th. 19] by Matthews and Wehner, and identifies the weakness of [10, Th. 1] by Wang and Renner in classical-quantum channel coding. For more general channels and entanglement-assisted codes, it is not clear whether the bounds in [11, Th. 18 and Th. 19] coincide with the exact error probability. To study this, a generalization of Theorem 1 imposing less structure over the test alternatives is needed. Theorem 2 shows that the minimum error probability can be written as an optimization problem involving information-spectrum measures. In particular, this expression shows that the Hayashi-Nagaoka lemma [13, Lemma 4] yields the exact error probability after optimizantion over its free parameters.

Acknowledgment

The problem studied here was suggested to the author by Alfonso Martinez. The author thanks him, Albert Guillèn i Fàbregas and William Matthews for stimulating discussions related to this work.

References

[1] C. W. Helstrom, "Detection theory and quantum mechanics," *Inf. and Control*, vol. 10, no. 3, pp. 254–291, 1967.

[2] P. A. Bakut and S. S. Shchurov, "Optimal detection of a quantum signal," *Probl. Peredachi Inf.*, vol. 4, no. 1, pp. 77–82, 1968, (in Russian, English translation: *Probl. Inf. Transm.*, vol. 4, pp. 61-65, 1968).

[3] A. S. Holevo, "An analog of the theory of statistical decisions in noncommutative theory of probability," *Trudy Moskov. Mat. Obšč.*, vol. 26, pp. 133–149, 1972, (in Russian).

[4] ——, "Statistical decision theory for quantum systems," *J. Multivariate Anal. 3*, vol. 3, no. 4, pp. 337–394, 1973.

[5] H. P. Yuen, R. S. Kennedy, and M. Lax, "Optimum testing of multiple hypotheses in quantum detection theory," *IEEE Trans. Inf. Theory*, vol. 21, no. 2, pp. 125–134, Mar 1975.

[6] C. W. Helstrom, *Quantum Detection and Estimation Theory*. NY: Academic Press, 1976.

[7] C. E. Shannon, R. G. Gallager, and E. R. Berlekamp, "Lower bounds to error probability for coding on discrete memoryless channels. I," *Inf. Contr.*, vol. 10, no. 1, pp. 65–103, 1967.

[8] H. Nagaoka, "Strong converse theorems in quantum information theory," in *Proc. ERATO Conf. Quantum Inf. Science*, Tokyo, Japan, 2001, p. 33.

[9] Y. Polyanskiy, H. V. Poor, and S. Verdú, "Channel coding rate in the finite blocklength regime," *IEEE Trans. Inf. Theory*, vol. 56, no. 5, pp. 2307–2359, 2010.

[10] L. Wang and R. Renner, "One-shot classical-quantum capacity and hypothesis testing," *Phys. Rev. Lett.*, vol. 108, no. 20, p. 200501, 2012.

[11] W. Matthews and S. Wehner, "Finite blocklength converse bounds for quantum channels," *IEEE Trans. Inf. Theory*, vol. 60, no. 11, pp. 7317–7329, 2014.

[12] H. Nagaoka and M. Hayashi, "An information-spectrum approach to classical and quantum hypothesis testing for simple hypotheses," *IEEE Trans. Inf. Theory*, vol. 53, no. 2, pp. 534–549, 2007.

[13] M. Hayashi and H. Nagaoka, "General formulas for capacity of classical-quantum channels," *IEEE Trans. Inf. Theory*, vol. 49, no. 7, pp. 1753–1768, 2003.

[14] G. Vazquez-Vilar, A. Tauste Campo, A. Guillén i Fàbregas, and A. Martinez, "Bayesian M-ary hypothesis testing: The meta-converse and Verdú-Han bounds are tight," *IEEE Trans. Inf. Theory*, 2016, to appear. Preprint available at arXiv:1411.3292.

[15] A. Jenčová, "Quantum hypothesis testing and sufficient subalgebras," *Lett. Math. Phys.*, vol. 93, no. 1, pp. 15–27, 2010.